

Use of ATLANTIS Tierra 2.0 in Mapping the Biodiversity (Invertebrates and Bryophytes) of Caves in the Azorean Archipelago

Paulo A.V. Borges^{1,2,3}, Rosalina Gabriel³, Fernando Pereira^{1,2,3}, Enésima P. Mendonça³, Eva Sousa³

¹ “Os Montanheiros”, Rua da Rocha, 9700 Angra do Heroísmo, Terceira, Açores, Portugal.

² GESPEA – Grupo de Estudo do Património Espeleológico dos Açores.

³ Universidade dos Açores, Dep. Ciências Agrárias, CITA-A, 9700-851 Angra do Heroísmo, Terceira, Açores; pborges@mail.angra.uac.pt.

Abstract

In this contribution the software ATLANTIS Tierra 2.0 is described as a promising tool to be used in the conservation management of the animal and plant biodiversity of caves in Macaronesia. In the Azores, the importance of cave entrances to bryophytes is twofold: i) since these are particularly humid, sheltered habitats, they support a diverse assemblage of bryophyte species and circa 25% of the Azorean bryoflora is referred to this habitat and ii) species, either endemic or referred in the European red list due to their vulnerability (19 species) or rarity (13) find refuge there. Cave adapted arthropods are also diverse in the Azores and 21 endemic obligate

cave species were recorded. Generally these species have restricted distributions and some are known from only one cave. ATLANTIS Tierra 2.0 allows the mapping of the distribution of all species in a 500 x 500 m grid in a GIS interface. This allows an easy detection of species rich caves (hotspots) and facilitates the interpretation of spatial patterns of species distribution. For instance, predictive models of species distribution could be constructed using the distribution of lava flows or other environmental variables. Using this new tool we will be better equipped to answer the following questions: a) Where are the current “hotspot caves” of biodiversity in the Azores?; b) How many new caves need to be selected as specially protected areas in

order to conserve the rarest endemic taxa?; c) Is there congruence between the patterns of richness and distribution of invertebrates and bryophytes?; d) Are environmental variables good surrogates of species distributions?

Introduction

The study of Azorean cave fauna and flora only started in 1988 with two expeditions of “National Geographic” under the supervision of Pedro Oromí (Univ. de La Laguna) and Philippe Ashmole (Univ. de Edinburg) and with the support of the speleological Azorean group “Os Montanheiros” (see Oromí *et al.* 1990, González-Mancebo *et al.* 1991). After those two expeditions in 1988 and 1990, the University of the Azores and “Os Montanheiros” performed most of the biospeleological work in the Azores (see Borges & Oromí 1994, 2006, Gabriel & Dias 1994). In the Azores, the importance of cave entrances to bryophytes is twofold: i) since these are particularly humid, sheltered habitats, they support a diverse assemblage of bryophyte species and circa 25% of the Azorean bryoflora is referred to this habitat and ii) species, either endemic or referred in the European red list (ECCB 1995) due to their vulnerability (19 species) or rarity (13) find refuge there. Cave adapted arthropods are also diverse in the Azores and 21 endemic obligate cave species were recorded (Borges & Oromí 2006). Generally these species have restricted distributions and some are known from only one cave (Borges & Oromí 2006).

There is a general agreement among scientists that biodiversity is under assault on a global basis and that species are being lost at greatly enhanced rates due to human processes such as habitat loss and fragmentation, invasive species, pollution and global climate change




Figure 1. Entrance window of ATLANTIS Tierra 2.0, in which it is possible to observe eight possible entrance gateways, the most relevant being the taxonomic reports (“Taxonomía”), information about species (“Consulta de especies”) and data analysis (“Consulta de análisis”).

(Lawton & May 1995; Chapin et al. 2000). Moreover, some recent studies indicate that there are some concerns related with invasive species and the conservation of native biodiversity in the Azores (Silva & Smith 2004, Borges et al. 2006).

In this contribution, a new software, ATLANTIS Tierra 2.0, is described as a promising tool to be used in the conservation management of the animal and plant biodiversity of caves from the Azores.

ATLANTIS Tierra 2.0

Since 1998 the Government of the Canary Islands has been conducting an important project on biodiversity, Project BIOTA (see Izquierdo *et al.* 2001, 2004). A Visual Basic software, called ATLANTIS Tierra 2.0, was developed for biodiversity data storage. With this database it will be possible to gather detailed information about all species on the surveyed geographical areas of

interest. This software has several important tools, namely a taxonomic tool and a conservation management analysis tool (Fig. 1) that allows the calculation of species richness, their rarity or complementarity in all 500x500 m cells of a particular island or, in any special area in one island.

With this software all the information we could think of about a species (e.g. the cavernicolous ground-beetle *Trechus montanheiorum*) is available in clicking the **information about species** (“**Consulta de especies**”) window (see Fig. 2). In this window it is also possible to check the detailed distribution of the species in a 500 x 500 m scale (Fig. 3). With this tool we may also investigate the distribution of the species throughout time in asking for its distribution in different time intervals. To each signalized 500 x 500 m grid cell correspond a cave for which the species was signalized in the literature.

However, it is in the data analysis

facility that ATLANTIS Tierra 2.0 is more interesting in terms of its application in a conservation management study. As an example in Fig. 4 we see the species richness of the European Rare Bryophytes (ECCB 1995) in caves from Graciosa Island (Azores). The grid-cell with the highest number of species corresponds to the location of Furna do Enxofre, currently a volcanic pit protected by law and under the special management of the Government. In Fig. 4 we can see also the list of species in grid cell with the highest number of species and that list could be exported to another software (e.g. Excel).

Very important in conservation management studies is to ask: “How many sites are needed to include all species of interest at least once?”. To answer this question, we could use the complementarity procedure, in which we get the minimum set of caves that combined have the highest representation of species (see Williams 2001). ATLANTIS


Consulta / Selección de Especies / Subespecies

Consulta de Especies/Subespecies

Código: A00017 Nombre común:

Validación taxón: Taxón válido Dcto. validación: A00002

Nombre especie: Trechus montanheiorum Oromí & Borges, 1991

Filo: Arthropoda Imagen básica: 

Clase: Insecta Más imágenes:

Orden: Coleoptera

Familia: Carabidae

Descripción:

Datos adicionales:

-
-
-
-
-

Endemismo del taxón seleccionado:

	Niv.territorial	Endémico	Dcto.Asig.
Azores			
Género:	—		
Especie:	Sí		A00002
Subespecie:	—		
Macaronesia			
Género:	—		
Especie:	—		
Subespecie:	—		

Registro << < 3 > >> de 9

Figure 2. Species management window of ATLANTIS Tierra 2.0, in which it is possible to observe the nomenclature of the species, a picture, the distribution of the species in the archipelago (green island) and other relevant information concerning the habitats, conservation status, biogeographical origin, etc.




Figure 3. Species management window of ATLANTIS Tierra 2.0, in which it is possible to observe the detailed distribution of *Trechus montanheiorum* in the island of Pico (lines are main roads in the island).




Figure 4. Data analysis window of ATLANTIS Tierra 2.0, in which it is possible to observe the number of bryophyte species in the European Red List present in caves from Graciosa Island (Azores). The list of species in the window corresponds to the grid cell with 15 species (Furna do Enxofre).




Figure 5. Data analysis window of ATLANTIS Tierra 2.0, in which it is possible to observe the four grid-cells that are necessary to include all the endemic arthropods occurring in caves from Terceira island (see text for further explanations).




Figure 6. Data analysis window of ATLANTIS Tierra 2.0, in which it is possible to observe the list of endemic arthropods that occur in two distinct cave systems at Terceira island (see text for further explanations).

Tierra 2.0 uses the heuristic suboptimal simple-greedy reserve-selection algorithm: first, the grid-cell with the highest species richness is selected. Then, these species are ignored and the grid-cell with the highest complement of species (that is, the most species not represented in the previous selected grid-cell), and so on, until all species are represented at least once. One good example of the application of the complementarity procedure is showed in Fig. 5, in which only four out of the eleven grid-cells with caves are necessary to protect the 26 endemic arthropod species occurring in the caves of this island. Those four grid-cells are signalized with a green dark border (the first selected grid-cell) and with a reddish dark border (the three other selected grid-cells). Therefore, with only four caves well managed we may protect all the endemic arthropod species known to occur in caves at Terceira Island (Azores). However, we should call attention to the fact that the complementarity procedure could be made more complex asking for the minimum set of caves that combined have at least each species represented twice, therefore assuring that species are protected in more than one place.

Another important facility available in ATLANTIS Tierra 2.0 is related with the investigation of the species composition in different areas of a region. For instance, we could have the list of species that are common in two different cave systems (Fig. 6). We could also get the list of species for each cave system and by exclusion obtain the lists of species that are exclusive to each cave system.

Conclusion

There is some urgency in the conservation of the diverse community of mosses and liverworts (Bryophyta) as well as of the rich cave adapted arthropods occurring in the Azorean lava tubes and

volcanic pits. The general pattern that emerges is that ATLANTIS Tierra 2.0 will be an important tool not only for the Azorean Government in managing the territory and designing natural protected areas, but also for research in the areas of applied ecology and conservation.

Using the ATLANTIS Tierra 2.0 new tool we will be better equipped to answer the following important questions: a) Where are the current “hotspot caves” of biodiversity in the Azores?; b) How many new caves need to be selected as specially protected areas in order to conserve the rarest endemic taxa?; c) Is there congruence between the patterns of richness and distribution of invertebrates and bryophytes?; d) Are environmental variables good surrogates of species distributions?

Acknowledgements

We wish to thank to Azorean Government for supporting our trip to Mexico to participate on the XIInd International Symposium on Vulcanospeleology (Tepoztlan, Morelos, México, July 2006).

Digital information of the islands was obtained within Project ATLÂNTICO—INTERREG IIIB, with permission of “Instituto Geográfico do Exército” under contract n° 047/CCO/2003.

References

- Borges, P.A.V., Lobo, J.M., Azevedo, E. B., Gaspar, C., Melo, C. & Nunes, L.V. (2006). Invasibility and species richness of island endemic arthropods: a general model of endemic vs. exotic species. *Journal of Biogeography* 33: 169-187.
- Borges, P.A.V. & Oromí, P. (1994). The Azores. In: C. Juberthie & V. Decu (Eds.) *Encyclopaedia Biospeleologica. Tome I*. pp. 605-610. Société de Biospéologie, Moulis.
- Borges, P.A.V. & Oromí, P. (2006). The

Azores. In: C. Juberthie & V. Decu (Eds.) *Encyclopaedia Biospeleologica. Tome Ia Amérique et Europe*. pp. ????. Société de Biospéologie, Moulis.

- Chapin III, F.S., Zavaleta, E.S., Eviner, V.T., Naylor, R.L., Vitousek, P.M., Reynolds, H.L., Hooper, D.U., Lavelle, S., Sala, O.E., Hobbie, S.E., Mack, M.V. & Díaz, S. (2000) Consequences of changing biodiversity. *Nature* 405: 234-242.
- ECCB (1995) *Red data book of European bryophytes*. European Committee for the Conservation of Bryophytes. Trondheim.
- Gabriel, R. & Dias, E. (1994). First approach to the study of the Algar do Carvão flora (Terceira, Azores). in: Actas do 3º Congresso Nacional de Espeleologia e do 1º Encontro Internacional de Vulcanospeleologia das Ilhas Atlânticas (30 de Setembro a 4 de Outubro de 1992), pp. 206-213. Angra do Heroísmo.
- González-Mancebo, J.M., Losada-Lima, A. & Hernández-García, C.D. (1991). A contribution to the floristic knowledge of caves on the Azores. *Mémoires de Biospéologie*, 18: 219-226.
- Lawton, J.H. & May, R.M. (1995) *Extinction Rates*. Oxford University Press, Oxford.
- Oromí, P., Martín, J.L., Ashmole, N.P. & Ashmole, M.J. (1990). A preliminary report on the cavernicolous fauna of the Azores. *Mémoires de Biospéologie*, 17: 97-105.
- Silva, L. & Smith, C.W. (2004) A characterization of the non-indigenous flora of the Azores Archipelago. *Biological Invasions*, 6: 193-204.
- Williams P. (2001). Complementarity. In: Levin S. (ed.), *Encyclopaedia of Biodiversity*, Volume 5. Academic Press, pp. 813-829.